

# Goat anti-Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 488

| Product Details         |                                            |
|-------------------------|--------------------------------------------|
| Size                    | 1 mg                                       |
| Species Reactivity      | Mouse                                      |
| Host/Isotype            | Goat / IgG                                 |
| Class                   | Polyclonal                                 |
| Type                    | Secondary Antibody                         |
| Conjugate               | Alexa Fluor™ 488                           |
| Excitation/Emission Max | 499/520 nm                                 |
| Immunogen               | Gamma Immunoglobins Heavy and Light chains |
| Form                    | Liquid                                     |
| Concentration           | 2 mg/mL                                    |
| Purification            | purified                                   |
| Storage buffer          | PBS, pH 7.5                                |
| Contains                | 5mM sodium azide                           |
| Storage conditions      | 4° C, store in dark                        |
| RRID                    | AB_2534069                                 |

| Applications                                      | Tested Dilution | Publications  |
|---------------------------------------------------|-----------------|---------------|
| Western Blot (WB)                                 | -               | 0 Publication |
| Immunohistochemistry (IHC)                        | Assay-dependent | 0 Publication |
| Immunohistochemistry (Paraffin) (IHC (P))         | -               | 0 Publication |
| Immunohistochemistry (PFA fixed) (IHC (PFA))      | -               | 0 Publication |
| Immunohistochemistry (Frozen) (IHC (F))           | -               | 0 Publication |
| Immunohistochemistry - Free Floating (IHC (Free)) | -               | 0 Publication |
| Immunocytochemistry (ICC/IF)                      | 1 µg/mL         | 0 Publication |
| Flow Cytometry (Flow)                             | 1-10 µg/mL      | 0 Publication |
| in situ PLA (PLA)                                 | -               | 0 Publication |
| Miscellaneous PubMed (Misc)                       | -               | 0 Publication |

## Product Specific Information

To minimize cross-reactivity, these goat anti-mouse IgG whole antibodies have been cross-adsorbed against human IgG and human serum. Cross-adsorption or pre-adsorption is a purification step to increase specificity of the antibody resulting in higher sensitivity and less background staining. The secondary antibody solution is passed through a column matrix containing immobilized serum proteins from potentially cross-reactive species. Only the nonspecific-binding secondary antibodies are captured in the column, and the highly specific secondaries flow through. The benefits of this extra step are apparent in multiplexing/multicolor-staining experiments (e.g., flow cytometry) where there is potential cross-reactivity with other primary antibodies or in tissue/cell fluorescent staining experiments where there may be the presence of endogenous immunoglobulins. For a highly cross-adsorbed secondary antibody equivalent, please see product Cat. No. A11029.

Alexa Fluor dyes are among the most trusted fluorescent dyes available today. Invitrogen™ Alexa Fluor 488 dye is a bright, green-fluorescent dye with excitation ideally suited to the 488 nm laser line. For stable signal generation in imaging and flow cytometry, Alexa Fluor 488 dye is pH-insensitive over a wide molar range. Probes with high fluorescence quantum yield and high photostability allow detection of low-abundance biological structures with great sensitivity. Alexa Fluor 488 dye molecules can be attached to proteins at high molar ratios without significant self-quenching, enabling brighter conjugates and more sensitive detection. The degree of labeling for each conjugate is typically 2-8 fluorophore molecules per IgG molecule; the exact degree of labeling is indicated on the certificate of analysis for each product lot.


Using conjugate solutions: Centrifuge the protein conjugate solution briefly in a microcentrifuge before use; add only the supernatant to the experiment. This step will help eliminate any protein aggregates that may have formed during storage, thereby reducing nonspecific background staining. Because staining protocols vary with application, the appropriate dilution of antibody should be determined empirically. For the fluorophore-labeled antibodies a final concentration of 1-10 µg/mL should be satisfactory for most immunohistochemistry and flow cytometry applications.


Product will be shipped at Room Temperature.

## Product Images For Goat anti-Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 488


### Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody (A-11001) in IHC

Simultaneous detection of expression of five genes in a whole-mount *Drosophila* embryo by fluorescence in situ hybridization (FISH) with five RNA probes. Red: sog labeled using aminoallyl UTP (Product # A21663, A32765) and Alexa Fluor® 647 succinimidyl ester (Product # A-20006, A20106). Green: ind labeled with DNP, followed by rabbit anti-dinitrophenyl-KLH IgG antibody (Product # A-6430) prelabeled with the Zenon® Alexa Fluor® 555 Rabbit IgG Labeling Kit (Product # Z-25305). Blue: en labeled with biotin and detected with HRP-streptavidin and Alexa Fluor® 405 tyramide (TSA™ Kit 39, Product # T30952). Yellow: wg labeled with digoxigenin and detected with sheep anti-digoxigenin IgG antibody and Alexa Fluor® 594 Donkey Anti-Sheep IgG antibody (Product # A-11016). Magenta: msh labeled with fluorescein and detected with mouse anti-fluorescein /Oregon Green® IgG<sub>2a</sub> antibody (Product # A-6421) and Alexa Fluor® 488 Goat Anti-Mouse IgG antibody (Product # A-11001, A11029). Image contributed by Dave Kosman and Ethan Bier, University of California, San Diego.





**Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody (A-11001) in ICC/IF**  
Microtubules of bovine pulmonary artery endothelial cells tagged with mouse monoclonal anti- $\alpha$ -tubulin antibody (Product # A11126) and subsequently probed with: Alexa Fluor® 488 Goat Anti-Mouse IgG antibody (Product # A-11001, top panel), Alexa Fluor® 546 Goat Anti-Mouse IgG antibody (Product # A-11003, middle panel) or Alexa Fluor® 594 Goat Anti-Mouse IgG antibody (Product # A-11005, bottom panel). These images were acquired using a FITC bandpass optical filter set, a rhodamine bandpass optical filter set, and a Texas Red bandpass optical filter set, respectively.



**Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody (A-11001) in IHC**  
Formation of the cephalic furrow in the anterior end of a developing *Drosophila melanogaster* embryo visualized with the help of several fluorescent stains. A primary antibody to neurotactin was visualized using a red-fluorescent Cy3 dye secondary antibody (Amersham Pharmacia Biotech Ltd.). Primary antibodies to plasma membrane-bound myosin and to nuclear-localized even-skipped (Eve) protein were visualized with green-fluorescent Alexa Fluor® 488 Goat Anti-Mouse IgG antibody (Product # A-11001) and Alexa Fluor® 488 Goat Anti-Rabbit IgG antibody (Product # A-11008), respectively. The nuclei were stained with blue-fluorescent Hoechst 33342 (Product # H1399, H3570, H21492). The sample was prepared by Eric Wieschaus, and the imaging was performed by Joe Goodhouse of Princeton University.

[View more figures on thermofisher.cn](http://thermofisher.cn)

Mouse strain-specific responses along the gut-brain axis upon fecal microbiota transplantation from children with autism. Gut Microbes (2025)

Ce6-GFFY is a novel photosensitizer for colorectal cancer therapy. Genes Dis (2025)

A Screen of Plant-Based Natural Products Revealed That Quercetin Prevents Pyroglutamylated Amyloid-(A3(pE)-42) Uptake in Astrocytes As Well As Resulting Astrogliosis and Synaptic Dysfunction. Mol Neurobiol (2025)

Nuclear talin-1 provides a bridge between cell adhesion and gene expression. iScience (2025)

Regeneration in the absence of canonical neoblasts in an early branching flatworm. Nat Commun (2025)

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLIED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCT(S) AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses, or any type of consumption by or application to human or animals.