



Goat anti-Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 568

Product Details	
Size	1 mg
Species Reactivity	Mouse
Host/Isotype	Goat / IgG
Class	Polyclonal
Туре	Secondary Antibody
Conjugate	Alexa Fluor™ 568
Excitation/Emission Max	579/603 nm
Immunogen	Gamma Immunoglobins Heavy and Light chains
Form	Liquid
Concentration	2 mg/mL
Purification	purified
Storage buffer	PBS, pH 7.5
Contains	5mM sodium azide
Storage conditions	4° C, store in dark
RRID	AB_144696

Applications	Tested Dilution	Publications
Western Blot (WB)	-	0 Publication
Immunohistochemistry (IHC)	-	0 Publication
Immunohistochemistry (Paraffin) (IHC (P))	-	0 Publication
Immunohistochemistry (Frozen) (IHC (F))	-	0 Publication
Immunocytochemistry (ICC/IF)	2 μg/mL	0 Publication
Flow Cytometry (Flow)	1-10 μg/mL	-
Miscellaneous PubMed (Misc)	-	0 Publication

Product Specific Information

To minimize cross-reactivity, these goat anti-mouse IgG (H+L) whole secondary antibodies have been affinity purified and cross-adsorbed against bovine IgG, goat IgG, rabbit IgG, rat IgG, human IgG, and human serum. Cross-adsorption or preadsorption is a purification step to increase specificity of the antibody resulting in higher sensitivity and less background staining. The secondary antibody solution is passed through a column matrix containing immobilized serum proteins from potentially cross-reactive species. Only the nonspecific-binding secondary antibodies are captured in the column, and the highly specific secondaries flow through. The benefits of this extra step are apparent in multiplexing/multicolor-staining experiments (e.g., flow cytometry) where there is potential cross-reactivity with other primary antibodies or in tissue/cell fluorescent staining experiments where there are may be the presence of endogenous immunoglobulins.

Alexa Fluor dyes are among the most trusted fluorescent dyes available today. Invitrogen™ Alexa Fluor 568 dye is a bright, orange/red-fluorescent dye with excitation ideally suited to the 568 nm laser line. For stable signal generation in imaging and flow cytometry, Alexa Fluor 568 dye is pH-insensitive over a wide molar range. Probes with high fluorescence quantum yield and high photostability allow detection of low-abundance biological structures with great sensitivity. Alexa Fluor 568 dye molecules can be attached to proteins at high molar ratios without significant self-quenching, enabling brighter conjugates and more sensitive detection. The degree of labeling for each conjugate is typically 2-8 fluorophore molecules per IgG molecule; the exact degree of labeling is indicated on the certificate of analysis for each product lot.

Using conjugate solutions: Centrifuge the protein conjugate solution briefly in a microcentrifuge before use; add only the supernatant to the experiment. This step will help eliminate any protein aggregates that may have formed during storage, thereby reducing nonspecific background staining. Because staining protocols vary with application, the appropriate dilution of antibody should be determined empirically. For the fluorophore-labeled antibodies a final concentration of 1-10 μ g/mL should be satisfactory for most immunohistochemistry and flow cytometry applications.

Product will be shipped at Room Temperature.

Product Images For Goat anti-Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 568


a b c c f

Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody (A-11031) in ICC/IF

Immunofluorescence analysis of Goat anti-Mouse IgG (H+L) Secondary Antibody, Alexa Fluor 568 conjugate (Product # A11031) was performed using HeLa cells stained with alpha Tubulin (236-10501) Mouse Monoclonal Antibody (Product # A11126). The cells were fixed with 4% paraformaldehyde for 10 minutes, permeabilized with 0.1% TritonTM X-100 for 10 minutes, blocked with 1% BSA for 1 hour and labeled with mouse primary antibody (1:250 dilution) for 3 hours at room temperature. Goat anti-Mouse IgG (H+L) Secondary Antibody, Alexa Fluor 568 conjugate was used at concentration of 2 μg/mL in phosphate buffered saline containing 0.2 % BSA for 45 minutes at room temperature, for detection of alpha Tubulin in the cytoplasm (Panel a: red). Nuclei (Panel b: blue) were stained with DAPI in SlowFade® Gold Antifade Mountant (Product # S36938). F-actin was stained with Alexa Fluor® 488 Phalloidin (Product # A12379, 1:300) (Panel c: green). Panel d represents the composite image. No nonspecific staining was observed with the secondary antibody alone (panel f), or with an isotype control (panel e). The images were captured at 60X magnification.


Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody (A-11031) in ICC/IF

MHCII+ adipocytes are functionally increased in cold-adapted iWAT. Eight-weekold male C57BL/6J mice were either housed at TN (30 °C) or put to cold temperature (18 °C for 14 days followed by 6 °C for 7 days). The mature adipocytes were isolated from the iWAT for further analysis. (A) mRNA expression of MHC-II associated genes and (B) immunofluorescence staining of HLA-DR in mature adipocytes. Scale bars, 200 µm. (C) Quantification of (B). (D) MLR assay to quantify the T cell proliferation in cocultures. (E) Subclustering of ASPCs. (F) RNA velocity of ASPCs and adipocyte subclusters. (G) Feature plots of Dpp4, Icam1, and Cd74 in ASPC subclusters. (H) Bubble chart of ASPC subclusters. (I) Percentage of CD74+ ASC in iWAT by flow cytometry. (J and K) Subcluster-10 and the rest of the ASPCs (control) were sorted from iWAT SVF and differentiated in vitro. (J) Adipogenesis was assessed by BODIPY staining. Scale bar, 200µm. (K) Relative mRNA expression of MHCII-related genes. (L) MLR assay to quantify the T cell proliferation in cocultures. *P < 0.05 and **P < 0.01. - Image collected and cropped by CiteAb under a CC-BY license from the following publication: Cold-Induced Reprogramming of Subcutaneous White Adipose Tissue Assessed by Single-Cell and Single-Nucleus RNA Sequencing. <i>Research (Wash D C)</i> (2023) Image collected and cropped by CiteAb from the following publication (https://pubmed.ncbi.nlm.nih.gov/37398933), licensed under a CC BY license.



Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody (A-11031) in ICC/IF

Loss of CAF-1 suppresses genomic instability in BRCA-deficient cells.a, b H2AX immunofluorescence experiment showing that CHAF1A depletion suppresses CPT-induced DNA damage accumulation in BRCA1 or BRCA2-depleted cells. HeLa cells were treated with 1 µM CPT for 1 h followed by media removal and chase in fresh media for 3 h. Representative micrographs (scale bar represents 50 µm) (a) and quantifications (b) are shown. At least 50 cells were quantified for each condition. The mean values are represented on the graph, and the p-values (t-test, two-tailed, unpaired) are listed at the top. Western blots confirming the codepletions are shown in Supplementary Fig. 2a. c-e Neutral comet assay showing that CHAF1A depletion suppresses CPT-induced (c, d) and cisplatininduced (e) DSB formation in BRCA1 or BRCA2-depleted cells. HeLa cells were treated with 100 nM CPT for 4 h or 2.5 µM cisplatin for 24 h. Representative micrographs (c) and quantifications (d, e) are shown. At least 60 nuclei were quantified for each condition. The mean values are represented on the graph, and the p-values (t-test, two-tailed, unpaired) are listed at the top. f, g Clonogenic survival experiments showing that CHAF1A co-depletion in BRCA2-knockdown (f) or BRCA1-knockdown (g) HeLa cells promotes cisplatin resistance. The average of three experiments, with standard deviations indicated as error bars, is shown. Asterisks indicate sta... Image collected and cropped by CiteAb from the following publication (https://pubmed.ncbi.nlm.nih.gov/36085347), licensed under a CC BY license.



View more figures on thermofisher.cn

□ 1725 References

Gross anatomy of the visual processing centers of Hieroglyphus banian. Cell Tissue Res (2025)

Structural, molecular and developmental evidence for cell-type diversity in cnidarian mechanosensory neurons. Nat Commun (2025)

Functional Characterization of Parallel Fiber-Purkinje Cell Synapses in Two Friedreich's Ataxia Mouse Models. Cerebellum (2025)

Elucidating subcellular architecture and dynamics at isotropic 100-nm resolution with 4Pi-SIM. Nat Methods (2025)

Driver lines for studying associative learning in Drosophila. Elife (2025)

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warneted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production occumentation; beginning package insents ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty immitted to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT.

BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED, ACCEDING TO FOR PEAIR, REPLACE OR REFUND FOR THE NON-CONFORMING PRODUCTS), AT SELLER'S SOLE OPTION, THERE IS NO A BUYER'S EXCLUSIVE THE WARRANTY PERIOD IS LIMITED. AND A SECRET OF THE PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, or vivo or in vivo therapeutic uses, or any type of consumption to human or animals.