

Goat anti-Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 790

Product Details	
Size	1 mg
Species Reactivity	Mouse
Host/Isotype	Goat / IgG
Class	Polyclonal
Type	Secondary Antibody
Conjugate	Alexa Fluor™ 790
Excitation/Emission Max	782/805 nm
Immunogen	Gamma Immunoglobins Heavy and Light chains
Form	Liquid
Concentration	2 mg/mL
Purification	purified
Storage buffer	PBS, pH 7.5
Contains	5mM sodium azide
Storage conditions	4° C, store in dark
RRID	AB_2534140


Applications	Tested Dilution	Publications
Western Blot (WB)	1:5,000-1:20,000	0 Publication
Immunocytochemistry (ICC/IF)	1-10 µg/mL	-
Flow Cytometry (Flow)	1-10 µg/mL	-

Product Specific Information


This secondary antibody is designed for fluorescent Western blot detection on various near-infrared fluorescence instruments. This antibody can be used for multi-color and multiplexing detection when using other antibodies conjugated to compatible Alexa Fluor™ dyes and wavelengths. Other applications of this antibody include immunofluorescent and fluorescent imaging applications when using instrumentation with appropriate excitation and detection capabilities.

Product will be shipped at Room Temperature.

Product Images For Goat anti-Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 790



Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody (A11357)
Specificity of secondary antibody was demonstrated by specific detection of the target immunoglobulin. Antibody specificity was demonstrated by specific detection of Mouse IgG. A band at ~55 kDa corresponding to Mouse IgG Heavy Chain was observed in Mouse IgG, Mouse IgG1, Mouse IgG2a, Mouse IgG2b and Mouse IgG3 but not in other species using Goat anti-Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 790 (Product # A11357) in Western Blot. {RE}



Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody (A11357) in ICC/IF

Immunofluorescence analysis of Goat anti-Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 790 (Product # A11357) was performed using SH-SY5Y (positive model) and HeLa (negative model) cells stained with Nestin Monoclonal Antibody (10C2), eBioscience™ (Product # 14-9843-80). The cells were fixed with 4% paraformaldehyde for 10 minutes, permeabilized with 0.1% Triton™ X-100 for 15 minutes, blocked with 2% BSA for 1 hour and labeled with 1:500 dilution of primary antibody overnight at 4C. Goat anti-Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 790 (Product # A11357, 1:2000 dilution) in 0.1% BSA in PBS for 1 hour at room temperature, was used for detection of Nestin in the cytoskeleton (Panel a: red). Nuclei (Panel b: blue) were stained with Hoechst33342 (Product # H1399). F-actin was stained with Alexa Fluor® 488 Phalloidin (Product # A12379, 1:500) (Panel c: green). Panel d represents the composite image. The specificity of the secondary antibody was proved by the absence of signal in HeLa (negative model for Nestin) due to no primary antibody binding (Panel e). Non-specific staining was not observed with secondary antibody alone (panel f). The images were captured at 40X magnification in CellInsight CX7 LZR High-Content Screening (HCS) Platform (Product # CX7A1110LZR) and externally deconvoluted (D.Sage et al./Methods 115 (2017) 28–41).

Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody (A11357) in WB

Multiplexed fluorescent western blot was performed using Goat anti-Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor 790 (Product # A11357). Whole cell extracts of A-431 (Lane 1, 2), HaCaT (Lane 3, 4, 5), SH-SY5Y (Lane 6, 7, 8), HeLa (Lane 9, 10) and MCF7 (Lane 11) were electrophoresed usingNuPAGE™ 4-12% Bis-Tris Protein Gel (Product # NP03222BOX). Resolved proteins were transferred onto nitrocellulose membrane (Product # IB23001) byiBlot® 2 Dry BlottingSystem (Product # IB21001). The blot was probed with Cytokeratin 5 Monoclonal Antibody (3E2F1) (Product # MA5-15347), Vimentin Rabbit IgG Polyclonal Antibody (Product # PA5-27231) and alpha Tubulin Monoclonal Antibody (YL1/2) (Product # MA1-80017). Secondary antibodies (Product # A-11357, 1:20,000), (Product # A-31577, 1:10,000) and (Product # A48269, 1:10,000) were used for detection of Cytokeratin 5, Vimentin and alpha Tubulin respectively. Fluorescent detection was performed usingiBright™FL1500 (Product # A44115). The anti-mouse secondary antibody (Product # A11357) specifically detects the mouse primary antibody.

View more figures on thermofisher.cn

26 References

Tryptophan As a New Member of RNA-Induced Silencing Complexes Prevents Colon Cancer Liver Metastasis. Adv Sci (Weinh) (2024)

The activation of LBH-CRYAB signaling promotes cardiac protection against I/R injury by inhibiting apoptosis and ferroptosis. iScience (2024)

Systemically administered wound-homing peptide accelerates wound healing by modulating syndecan-4 function. Nat Commun (2023)

MYC disrupts transcriptional and metabolic circadian oscillations in cancer and promotes enhanced biosynthesis. PLoS Genet (2023)

Cardiac Protein Kinase D1 ablation alters the myocytes -adrenergic response. J Mol Cell Cardiol (2023)

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLIED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON-INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCT(S) AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses, or any type of consumption by or application to human or animals.