

**PRODUCT INFORMATION****SchI (MlyI)****#ER1371** 1000 U**Lot:** \_\_\_\_\_ **Expiry Date:** \_\_\_\_\_

5'...G A G T C (N)<sub>5</sub>↓...3'  
3'...C T C A G (N)<sub>5</sub>↑...5'

Concentration: 10 U/µL

Source: *E.coli* that carries the cloned *schlR* gene  
from *Staphylococcus cohnii* Lki 19-320

Supplied with: 1 mL of 10X Buffer Tango

**Store at -20°C**

BSA included

[www.thermoscientific.com/onebio](http://www.thermoscientific.com/onebio)**RECOMMENDATIONS****1X Thermo Scientific Tango Buffer** (for 100% SchI digestion)33 mM Tris-acetate (pH 7.9), 10 mM magnesium acetate,  
66 mM potassium acetate, 0.1 mg/mL BSA.**Incubation temperature**

37°C.

**Unit Definition**

One unit is defined as the amount of SchI required to digest 1 µg of lambda DNA in 1 hour at 37°C in 50 µL of recommended reaction buffer.

**Dilution**

Dilute with Dilution Buffer (#B19): 10 mM Tris-HCl (pH 7.4 at 25°C), 100 mM KCl, 1 mM EDTA, 1 mM DTT, 0.2 mg/mL BSA and 50% glycerol.

**Double Digests**Tango™ Buffer provided simplifies buffer selection for double digests. 98% of Thermo Scientific restriction enzymes are active in a 1X or 2X concentration of Tango Buffer. Please go to [www.thermoscientific.com/doubledigest](http://www.thermoscientific.com/doubledigest) to choose the best buffer for your experiments.**Storage Buffer**

SchI is supplied in: 10 mM Tris-HCl (pH 7.5 at 25°C), 100 mM KCl, 0.1 mM EDTA, 1 mM DTT, 0.2 mg/mL BSA and 50% glycerol.

## Recommended Protocol for Digestion

- Add:

nuclease-free water	16 $\mu$ L
10X Buffer Tango	2 $\mu$ L
DNA (0.5-1 $\mu$ g/ $\mu$ L)	1 $\mu$ L
Schl	0.5-2 $\mu$ L*
- Mix gently and spin down for a few seconds.
- Incubate at 37°C for 1-16 hours\*.

The digestion reaction may be scaled either up or down.

## Recommended Protocol for Digestion of PCR Products Directly after Amplification

- Add:

PCR reaction mixture	10 $\mu$ L (~0.1-0.5 $\mu$ g of DNA)
nuclease-free water	18 $\mu$ L
10X Buffer Tango	2 $\mu$ L
Schl	1-2 $\mu$ L*
- Mix gently and spin down for a few seconds.
- Incubate at 37°C for 1-16 hours\*.

\* See Star activity.

## Thermal Inactivation

Schl is inactivated by incubation at 65°C for 20 min.

## ENZYME PROPERTIES

### Enzyme Activity in Thermo Scientific REase Buffers, %

B	G	O	R	Tango	2X Tango
20-50	50-100	0-20	0-20	100	0-20

### Star Activity

An excess of Schl (15 U/ $\mu$ g DNA x 1 hour) may result in star activity.

### Methylation Effects on Digestion

Dam: never overlaps – no effect.  
Dcm: never overlaps – no effect.  
CpG: may overlap – no effect.  
EcoKI: never overlaps – no effect.  
EcoBI: may overlap – effect not determined.

### Stability during Prolonged Incubation

A minimum of 0.2 units of the enzyme is required for complete digestion of 1  $\mu$ g of lambda DNA in 16 hours at 37°C.

### Number of Recognition Sites in DNA

$\lambda$	$\Phi$ X174	pBR322	pUC57	pUC18/19	pTZ19R/U	M13mp18/19
61	10	4	3	4	7	8

### Note

Schl may remain associated with the cleaved DNA. This may cause DNA band shifting during electrophoresis. To avoid atypical DNA band patterns, use the 6X DNA Loading Dye&SDS Solution (#R1151) for sample preparation or heat the digested DNA in the presence of SDS prior to electrophoresis.

For **CERTIFICATE OF ANALYSIS** see back page

# CERTIFICATE OF ANALYSIS

## Overdigestion Assay

No detectable change in the specific fragmentation pattern is observed after a 10-fold overdigestion with SchI (10 U/ $\mu$ g lambda DNA x 1 hour) (see Star activity).

## Ligation and Recleavage (L/R) Assay

The ligation and reclavage assay was replaced with LO test after validating experiments showed LO test ability to trace nuclease and phosphatase activities with sensitivity that is higher than L/R by a factor of 100.

## Labeled Oligonucleotide (LO) Assay

No detectable degradation of single-stranded or double-stranded labeled oligonucleotides occurred during incubation with 10 units of SchI for 4 hours.

Quality authorized by:



Jurgita Zilinskiene

## PRODUCT USE LIMITATION

This product is developed, designed and sold exclusively *for research purposes and in vitro use only*. The product was not tested for use in diagnostics or for drug development, nor is it suitable for administration to humans or animals.

Please refer to [www.thermoscientific.com/onebio](http://www.thermoscientific.com/onebio) for Material Safety Data Sheet of the product.

© 2012 Thermo Fisher Scientific Inc. All rights reserved. All trademarks are the property of Thermo Fisher Scientific Inc. and its subsidiaries.

