

PRODUCT INFORMATION**SatI (Fnu4HI)****#ER1641** 200 U

Lot: _____ Expiry Date: _____

5'...G C↓N G C...3'

3'...C G N↑C G...5'

Concentration: 10 U/μL

Source: *Staphylococcus arlettae* RFL832

Supplied with: 1 mL of 10X Buffer G

1 mL of 10X Buffer Tango

Store at -20°C

BSA included

www.thermoscientific.com/onebio**RECOMMENDATIONS****1X Buffer G** (for 100% SatI digestion)10 mM Tris-HCl (pH 7.5), 10 mM MgCl₂, 50 mM NaCl, 0.1 mg/mL BSA.**Incubation temperature**

37°C.

Unit Definition

One unit is defined as the amount of SatI required to digest 1 μg of lambda DNA in 1 hour at 37°C in 50 μL of recommended reaction buffer.

Dilution

Dilute with Dilution Buffer (#B19): 10 mM Tris-HCl (pH 7.4 at 25°C) 100 mM KCl, 1 mM EDTA, 1 mM DTT, 0.2 mg/mL BSA and 50% glycerol.

Double Digests

Thermo Scientific Tango Buffer is provided to simplify buffer selection for double digests. 98% of Thermo Scientific restriction enzymes are active in a 1X or 2X concentration of Tango™ Buffer. Please refer to www.thermoscientific.com/doubledigest to choose the best buffer for your experiments.

1X Tango Buffer: 33 mM Tris-acetate (pH 7.9 at 37°C), 10 mM magnesium acetate, 66 mM potassium acetate, 0.1 mg/mL BSA.

Storage Buffer

SatI is supplied in: 10 mM Tris-HCl (pH 7.4 at 25°C), 100 mM KCl, 1 mM EDTA, 1 mM DTT, 0.2 mg/mL BSA and 50% glycerol.

Recommended Protocol for Digestion

- Add:

| | |
|------------------------------|---------------|
| nuclease-free water | 16 μ L |
| 10X Buffer G | 2 μ L |
| DNA (0.5-1 μ g/ μ L) | 1 μ L |
| SatI | 0.5-2 μ L |
- Mix gently and spin down for a few seconds.
- Incubate at 37°C for 1-16 hours.

The digestion reaction may be scaled either up or down.

Recommended Protocol for Digestion of PCR Products

Directly after Amplification

- Add:

| | |
|----------------------|--------------------------------------|
| PCR reaction mixture | 10 μ L (~0.1-0.5 μ g of DNA) |
| nuclease-free water | 18 μ L |
| 10X Buffer G | 2 μ L |
| SatI | 1-2 μ L |
- Mix gently and spin down for a few seconds.
- Incubate at 37°C for 1-16 hours.

Thermal Inactivation

SatI is inactivated by incubation at 65°C for 20 min.

ENZYME PROPERTIES

Enzyme Activity in Thermo Scientific REase Buffers, %

| B | G | O | R | Tango | 2X Tango |
|-------|-----|-------|-------|--------|----------|
| 20-50 | 100 | 20-50 | 20-50 | 50-100 | 20-50 |

Methylation Effects on Digestion

Dam: never overlaps – no effect.

Dcm: never overlaps – no effect.

CpG: may overlap – blocked.

EcoKI: never overlaps – no effect.

EcoBI: never overlaps – no effect.

Stability during Prolonged Incubation

A minimum of 0.1 units of the enzyme is required for complete digestion of 1 μ g of lambda DNA in 16 hours at 37°C.

Compatible Ends

GC↓(C/G)GC – Bcnl, Bme1390I

GC↓(A/T)GC – Mval, Bme1390I

Number of Recognition Sites in DNA

| λ | Φ X174 | pBR322 | pUC57 | pUC18/19 | pTZ19R/U | M13mp18/19 |
|-----------|-------------|--------|-------|----------|----------|------------|
| 380 | 31 | 42 | 19 | 19 | 20 | 17 |

Note

At least two copies of SatI recognition site are required for efficient cleavage.

For **CERTIFICATE OF ANALYSIS** see back page

CERTIFICATE OF ANALYSIS

Overdigestion Assay

No detectable change in the specific fragmentation pattern is observed after a 160-fold overdigestion with SstI (10 U/μg lambda DNA x 16 hours).

Ligation and Recleavage (L/R) Assay

The ligation and reclavage assay was replaced with LO test after validating experiments showed LO test ability to trace nuclease and phosphatase activities with sensitivity that is higher than L/R by a factor of 100.

Labeled Oligonucleotide (LO) Assay

No detectable degradation of single-stranded or double-stranded labeled oligonucleotides occurred during incubation with 10 units of SstI for 4 hours.

Quality authorized by:



Jurgita Zilinskiene

PRODUCT USE LIMITATION

This product is developed, designed and sold exclusively *for research purposes and in vitro use only*. The product was not tested for use in diagnostics or for drug development, nor is it suitable for administration to humans or animals.

Please refer to www.thermoscientific.com/onebio for Material Safety Data Sheet of the product.

© 2012 Thermo Fisher Scientific Inc. All rights reserved. All trademarks are the property of Thermo Fisher Scientific Inc. and its subsidiaries.