

## PRODUCT INFORMATION

# BshNI (BanI)

**#ER1001**      2000 U

**Lot:** \_\_\_\_      **Expiry Date:** \_\_

5'...**G↓G** Y R C C...3'

3'...C C R Y **G↑G**...5'

Concentration:      10 U/μL

Source:              *Bacillus sphaericus* Tk 4-5

Supplied with:      1 mL of 10X Buffer 0  
                                 1 mL of 10X Buffer Tango

**Store at -20°C**



BSA included

## RECOMMENDATIONS

**1X Buffer 0** (for 100% BshNI digestion)

50 mM Tris-HCl (pH 7.5), 10 mM MgCl<sub>2</sub>, 100 mM NaCl,  
0.1 mg/mL BSA.

**Incubation temperature**

37°C.

**Unit Definition**

One unit is defined as the amount of BshNI required to digest 1 μg of lambda DNA *dcm*<sup>-</sup> in 1 hour at 37°C in 50 μL of recommended reaction buffer.

**Dilution**

Dilute with Dilution Buffer (#B19): 10 mM Tris-HCl, (pH 7.4 at 25°C), 100 mM KCl, 1 mM EDTA, 1 mM DTT, 0.2 mg/mL BSA and 50% glycerol.

**Double Digests**

Thermo Scientific Tango Buffer is provided to simplify buffer selection for double digests. 98% of Thermo Scientific restriction enzymes are active in a 1X or 2X concentration of Tango™ Buffer. Please refer to [www.thermoscientific.com/doubledigest](http://www.thermoscientific.com/doubledigest) to choose the best buffer for your experiments.

1X Tango Buffer: 33 mM Tris-acetate (pH 7.9 at 37°C), 10 mM magnesium acetate, 66 mM potassium acetate, 0.1 mg/mL BSA.

## Storage Buffer

BshNI is supplied in: 10 mM Tris-HCl (pH 7.5 at 25°C), 200 mM KCl, 1 mM DTT, 0.1 mM EDTA, 0.2 mg/mL BSA and 50% glycerol.

## Recommended Protocol for Digestion

- Add:

nuclease-free water	16 $\mu$ L
10X Buffer O	2 $\mu$ L
DNA (0.5-1 $\mu$ g/ $\mu$ L)	1 $\mu$ L
BshNI	0.5-2 $\mu$ L
- Mix gently and spin down for a few seconds.
- Incubate at 37°C for 1-16 hours.

The digestion reaction may be scaled either up or down.

## Recommended Protocol for Digestion of PCR Products Directly after Amplification

- Add:

PCR reaction mixture	10 $\mu$ L (~0.1-0.5 $\mu$ g of DNA)
nuclease-free water	18 $\mu$ L
10X Buffer O	2 $\mu$ L
BshNI	1-2 $\mu$ L
- Mix gently and spin down for a few seconds.
- Incubate at 37°C for 1-16 hours.

## Thermal Inactivation

BshNI is inactivated by incubation at 65°C for 20 min.

## ENZYME PROPERTIES

### Enzyme Activity in Thermo Scientific REase Buffers, %

B	G	O	R	Tango	2X Tango
0-20	20-50	100	50-100	0-20	100

### Methylation Effects on Digestion

Dam: never overlaps – no effect.  
Dcm: may overlap – cleavage impaired.  
CpG: may overlap – cleavage impaired.  
EcoKI: may overlap – effect not determined.  
EcoBI: never overlaps – no effect.

### Stability during Prolonged Incubation

A minimum of 0.3 units of the enzyme is required for complete digestion of 1  $\mu$ g of lambda DNA in 16 hours at 37°C.

### Digestion of Agarose-embedded DNA

A minimum of 5 units of the enzyme are required for complete digestion of 1  $\mu$ g of agarose-embedded lambda DNA in 16 hours.

### Compatible Ends

G↓GCGCC - SspDI  
G↓GTACC - Acc65I, Bsp1407I, Pfl123II, TatI

### Number of Recognition Sites in DNA

$\lambda$	$\Phi$ X174	pBR322	pUC57	pUC18/19	pTZ19R/U	M13mp18/19
25	3	9	4	4	4	7

### Note

BshNI cleavage is impaired by overlapping *dcm* methylation. To avoid *dcm* methylation, use a *dam*<sup>-</sup>, *dcm*<sup>-</sup> strain such as GM2163 (#M0099).

For **CERTIFICATE OF ANALYSIS** see back page

# CERTIFICATE OF ANALYSIS

## Overdigestion Assay

No detectable change in the specific fragmentation pattern is observed after a 160-fold overdigestion with BshNI (10 U/ $\mu$ g lambda DNA *dcm*<sup>-</sup> x 16 hours).

## Ligation and Recleavage (L/R) Assay

The ligation and recleavage assay was replaced with LO test after validating experiments showed LO test ability to trace nuclease and phosphatase activities with sensitivity that is higher than L/R by a factor of 100.

## Labeled Oligonucleotide (LO) Assay

No detectable degradation of single-stranded or double-stranded labeled oligonucleotides occurred during incubation with 10 units of BshNI for 4 hours.

Quality authorized by:



Jurgita Zilinskiene

## **PRODUCT USE LIMITATION**

This product is developed, designed and sold exclusively *for research purposes and in vitro use only*. The product was not tested for use in diagnostics or for drug development, nor is it suitable for administration to humans or animals.

Please refer to [www.thermoscientific.com/onebio](http://www.thermoscientific.com/onebio) for Material Safety Data Sheet of the product.

© 2012 Thermo Fisher Scientific Inc. All rights reserved. All trademarks are the property of Thermo Fisher Scientific Inc. and its subsidiaries.

