

PRODUCT INFORMATION

Nb.Mva1269I (Nb.BsmI)

#ER2051 1000 U

Lot: _____ Expiry Date: __

5'...G A A T G C...3'

3'...C T T A C↑G...5'

Concentration: 10 U/μL

Source: *E.coli* that carries the mutagenized *mva1269IR* gene from *Micrococcus varians* RFL1269

Supplied with: 1 mL of 10X Buffer O
1 mL of 10X Buffer Tango

Store at -20°C



BSA included

www.thermoscientific.com/onebio

DESCRIPTION

Nb.Mva1269I is a site and strand specific endonuclease artificially engineered from R.Mva1269I that cleaves only one strand of the DNA within its recognition sequence on double stranded DNA substrate.

RECOMMENDATIONS

1X Buffer O (for 100% Nb.Mva1269I digestion)

50 mM Tris-HCl (pH 7.5), 10 mM MgCl₂, 100 mM NaCl, 0.1 mg/mL BSA.

Incubation temperature

37°C.

Unit Definition

One unit is defined as the amount of Nb. Mva1269I required to convert 1 μg of supercoiled pBR322 DNA to circular form in 1 hour at 37°C in 50 μL of recommended reaction buffer.

Dilution

Dilute with Dilution Buffer (#B19): 10 mM Tris-HCl (pH 7.4 at 25°C), 100 mM KCl, 1mM EDTA, 1 mM DTT, 0.2 mg/mL BSA and 50% glycerol.

Storage Buffer

Nb. Mva1269I is supplied in: 10 mM Tris-HCl (pH 7.4 at 25°C), 50 mM NaCl, 1 mM DTT, 1 mM EDTA, 0.2 mg/mL BSA and 50% glycerol.

Thermal Inactivation

Nb.Mva1269I is inactivated by incubation at 80°C for 20 min.

ENZYME PROPERTIES

Enzyme Activity in Thermo Scientific REase Buffers, %

B	G	O	R	Tango	2X Tango
0-20	20-50	100	100	20-50	50-100

Stability during Prolonged Incubation

A minimum of 0.3 units of the enzyme is required for complete conversion of 1 µg of pBR322 DNA in 16 hours at 37°C into circular form.

Number of Recognition Sites in DNA

λ	ΦX174	pBR322	pUC57	pUC18/19	pTZ19R/U	M13mp18/19
46	4	1	1	0	0	1

APPLICATIONS

- Production of single-stranded circular DNA from supercoiled double-stranded plasmids *in vitro* with subsequent use in DNA sequencing, site-specific mutagenesis, etc.
- Creation of nested deletions.
- Vector preparation for ligation independent cloning method.
- Preparations of covalently closed, double-stranded linear DNA molecules.

This product is licensed under U.S. Patent No. 7,081,358 or corresponding foreign patents.

PRODUCT USE LIMITATION

This product is developed, designed and sold exclusively *for research purposes and in vitro use only*. The product was not tested for use in diagnostics or for drug development, nor is it suitable for administration to humans or animals.

Please refer to www.thermoscientific.com/onebio for Material Safety Data Sheet of the product.

© 2012 Thermo Fisher Scientific Inc. All rights reserved. All trademarks are the property of Thermo Fisher Scientific Inc. and its subsidiaries.

CERTIFICATE OF ANALYSIS

Nonspecific Nicking Assay

Incubation of 10 units of Nb.Mva1269I with 1 µg pUC19 DNA (lacking the recognition sequence of Nb.Mva1269I) for 16 hours at 37°C in 50 µL reaction buffer resulted in <3% conversion to circular form.

Double Strand Endonuclease Assay

Incubation of 10 unit of Nb.Mva1269I with 1 µg pBR322 DNA for 16 hours at 37°C in 50 µL reaction buffer resulted in <1% conversion to linear form.

Labeled Oligonucleotide (LO) Assay

No detectable degradation of single-stranded or double-stranded labeled oligonucleotides occurred during incubation with 10 units of Nb.Mva1269I for 4 hours.

Blue/White (B/W) Cloning Assay

The B/W assay was replaced with LO test after validating experiments showed LO test ability to detect nuclease and phosphatase activities with sensitivity that equals to that of B/W test.

Quality authorized by:



Jurgita Zilinskiene