

## PRODUCT INFORMATION

# SsiI (AciI)

**#ER1791      200 U**

**Lot: \_\_\_\_\_      Expiry Date: \_\_\_\_\_**

5'...C↓C G C...3'

3'...G G C↑G...5'

Concentration: 10 U/μL  
Source: *Staphylococcus sciuri* RFL1  
Supplied with: 1 mL of 10X Buffer O  
1 mL of 10X Buffer Tango

**Store at -20°C**



## RECOMMENDATIONS

**1X Buffer O** (for 100% SsiI digestion)

50 mM Tris-HCl (pH 7.5), 10 mM MgCl<sub>2</sub>, 100 mM NaCl, 0.1 mg/mL BSA.

**Incubation temperature**

37 °C.

**Unit Definition**

One unit is defined as the amount of SsiI required to digest 1 μg of lambda DNA in 1 hour at 37°C in 50 μL of recommended reaction buffer.

**Dilution**

Dilute with Dilution Buffer (#B19): 10 mM Tris-HCl, (pH 7.4 at 25 °C), 100 mM KCl, 1 mM EDTA, 1 mM DTT, 0.2 mg/mL BSA and 50% glycerol.

**Double Digests**

Thermo Scientific Tango Buffer is provided to simplify buffer selection for double digests. 98% of Thermo Scientific restriction enzymes are active in a 1X or 2X concentration of Tango™ Buffer. Please refer to [www.thermoscientific.com/doubledigest](http://www.thermoscientific.com/doubledigest) to choose the best buffer for your experiments.

1X Tango Buffer: 33 mM Tris-acetate (pH 7.9 at 37 °C), 10 mM magnesium acetate, 66 mM potassium acetate, 0.1 mg/mL BSA.

## Storage Buffer

Ssil is supplied in: 10 mM Tris-HCl (pH 7.4 at 25 °C), 100 mM KCl, 1 mM EDTA, 1 mM DTT, 0.2 mg/mL BSA and 50% glycerol.

## Recommended Protocol for Digestion

- Add:

nuclease-free water	16 µL
10X Buffer O	2 µL
DNA (0.5-1 µg/µL)	1 µL
Ssil	0.5-2 µL*
- Mix gently and spin down for a few seconds.
- Incubate at 37° C for 1-16 hours\*.

The digestion reaction may be scaled either up or down.

## Recommended Protocol for Digestion of PCR Products Directly after Amplification

- Add:

PCR reaction mixture	10 µL	(~0.1-0.5 µg of DNA)
nuclease-free water	18 µL	
10X Buffer O	2 µL	
Ssil	1-2 µL*	
- Mix gently and spin down for a few seconds.
- Incubate at 37°C for 1-16 hours\*.

\* See Overdigestion Assay.

## Thermal Inactivation

Ssil is inactivated by incubation at 65 °C for 20 min.

## ENZYME PROPERTIES

### Enzyme Activity in Thermo Scientific REase Buffers, %

B	G	O	R	Tango	2X Tango
NR	20-50	100	50-100	NR	100

NR – buffer is not recommended, because of high star activity

### Methylation Effects

- Dam: never overlaps – no effect.
- Dcm: never overlaps – no effect.
- CpG: completely overlaps – blocked.
- EcoKI: never overlaps – no effect.
- EcoBI: never overlaps – no effect.

### Stability during Prolonged Incubation

A minimum of 0.5 units of the enzyme is required for complete digestion of 1 µg of lambda DNA in 16 hours at 37 °C.

### Compatible Ends

Bsp119I, Bsu15I, Hin1I (GR/CGCC), Hin6I, HpaII, MspI, NarI, Psp1406I, TaqI, XmiI (GT/CGAC).

### Number of Recognition Sites in DNA

λ	ΦX174	pBR322	pUC57	pUC18/19	pTZ19R/U	M13mp18/19
516	36	67	34	34	32	42

For **CERTIFICATE OF ANALYSIS** see back page

# CERTIFICATE OF ANALYSIS

## Overdigestion Assay

No detectable change in the specific fragmentation pattern is observed after an 80-fold overdigestion with Ssi I (5 U/ $\mu$ g control DNA\* x 16 hours).

\*The control DNA is pBluescript II KS (+)DNA with the inserted Ssi I (Acil) recognition site.

## Ligation and Recleavage (L/R) Assay

The ligation and recleavage assay was replaced with LO test after validating experiments showed LO test ability to trace nuclease and phosphatase activities with sensitivity that is higher than L/R by a factor of 100.

## Labeled Oligonucleotide (LO) Assay

No detectable degradation of single-stranded or double-stranded labeled oligonucleotides occurred during incubation with 10 units of Ssi I for 4 hours.

Quality authorized by:



Jurgita Zilinskiene

**LIMITED USE LABEL LICENSE: Internal Research and Development Use Only.**

The purchase of this product conveys to the buyer the limited, non-exclusive, non-transferable right (without the right to resell, repackage, or further sublicense) to use this product for internal research and development purposes. No other license is granted to the buyer whether expressly, by implication, by estoppel or otherwise. In particular, the purchase of the product does not include or carry any right or license to use, develop, or otherwise exploit this product commercially and no rights are conveyed to the buyer to use the product or components of the product for purposes including but not limited to provision of services to a third party, generation of commercial databases or clinical diagnostics. This product is sold pursuant to authorization from Thermo Fisher Scientific and Thermo Fisher Scientific reserves all other rights. For information on purchasing a license for uses other than internal research and development purposes, please contact [outlicensing@lifetech.com](mailto:outlicensing@lifetech.com) or Out Licensing, Life Technologies Inc., 5791 Van Allen Way, Carlsbad, California 92008.

**PRODUCT USE LIMITATION**

This product is developed, designed and sold *exclusively for research purposes and in vitro use only*. The product was not tested for use in diagnostics or for drug development, nor is it suitable for administration to humans or animals.

Please refer to [www.thermoscientific.com/onebio](http://www.thermoscientific.com/onebio) for Material Safety Data Sheet of the product.

© 2015 Thermo Fisher Scientific Inc. All rights reserved. All trademarks are the property of Thermo Fisher Scientific Inc. and its subsidiaries.