

Goat anti-Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 660


Product Details	
Size	1 mg
Species Reactivity	Mouse
Host/Isotype	Goat / IgG
Class	Polyclonal
Type	Secondary Antibody
Conjugate	Alexa Fluor™ 660
Excitation/Emission Max	663/691 nm
Immunogen	Gamma Immunoglobins Heavy and Light chains
Form	Liquid
Concentration	2 mg/mL
Purification	purified
Storage buffer	PBS, pH 7.5
Contains	5mM sodium azide
Storage conditions	4°C, store in dark
RRID	AB_2535721

Applications	Tested Dilution	Publications
Western Blot (WB)	1:10,000	-
Immunohistochemistry (Frozen) (IHC (F))	-	0 Publication
Immunocytochemistry (ICC/IF)	1-10 µg/mL	-
Miscellaneous PubMed (Misc)	-	0 Publication


Product Specific Information

Product will be shipped at Room Temperature.


Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody (A-21054) in ICC/IF
Immunofluorescence analysis of Goat anti-Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor 660 (Product # A-21054) was performed using SH-SY5Y (positive model) and T-47D (negative model) cells stained with Nestin Monoclonal Antibody (10C2), eBioscience™ (Product # 14-9843-80). The cells were fixed with 4% paraformaldehyde for 10 minutes, permeabilized with 0.1% Triton™ X-100 for 10 minutes, blocked with 1% BSA for 1 hour and labeled with 2 µg/mL primary antibody for 3 hours at room temperature. Goat anti-Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor 660 (Product # A-21054, 1:2000 dilution) in 0.1% BSA in PBS for 45 minutes at room temperature, was used for detection of Nestin in the cytoskeleton (Panel a: Pink). Nuclei (Panel b: blue) were stained with Hoechst33342 (Product # H1399). F-actin was stained with Alexa Fluor® 488 Phalloidin (Product # A12379, 1:300) (Panel c: green). Panel d represents the composite image. The specificity of the secondary antibody was proved by the absence of signal in T-47D (negative model for Nestin) due to no primary antibody binding (Panel e). Non-specific staining was not observed with secondary antibody alone (panel f). The images were captured at 40X magnification in CellInsight CX7 LZR High-Content Screening (HCS) Platform (Product # CX7A1110LZR) and externally deconvoluted (D.Sage et al. /Methods 115 (2017) 28–41).



Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody (A-21054) in WB
Multiplexed fluorescent western blot was performed using Goat anti-Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor 660 (Product # A-21054). Whole cell extracts of HeLa (Lane 1), Hep G2 (Lane 2), U-87 MG (Lane 3, 4, 5), SH-SY5Y (Lane 6, 7, 8) were electrophoresed using NuPAGE™ 4-12% Bis-Tris Protein Gel (Product # NP03222BOX). Resolved proteins were transferred onto a nitrocellulose membrane (Product # IB23001) by iBlot® 2 Dry Blotting System (Product # IB21001). The blot was probed with PGP9.5 Monoclonal Antibody (13C4) (Product # MA1-90008) and HSP70 Polyclonal Antibody (Product # PA5-28003). Secondary antibodies (Product # A-21054, 1:10000 dilution) and (Product # A32808, 1:20000 dilution) were used for detection of PGP9.5 and HSP70 respectively. Fluorescent detection was performed using iBright FL1500 (Product # A44115). Goat anti-Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor 660 (Product # A-21054) specifically detects the mouse primary antibody.



Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody (A-21054) in WB
Western blot was performed using Goat anti-Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor 660 (Product # A-21054) and ~50, 25 kDa bands corresponding to Mouse IgG Heavy and Light Chain was observed in Mouse IgG, Mouse IgG1, Mouse IgG2a, Mouse IgG2b and Mouse IgG3 but not in Mouse IgM, Rabbit IgG, Chicken IgY, Goat IgG, Human IgG. Purified protein (100 ng) of Mouse IgG (Lane 1), Mouse IgM (Lane 2), Rabbit IgG (Lane 3), Rat IgG (Lane 4), Chicken IgY (Lane 5), Goat IgG (Lane 6) Human IgG (Lane 7), Mouse IgG1 (Lane 8), Mouse IgG2a (Lane 9) and Mouse IgG3 (Lane 10) (Fig. a) were electrophoresed using NuPAGE™ 4-12% Bis-Tris Protein Gel (Product # NP03222BOX). Resolved proteins were then transferred onto a nitrocellulose membrane (Product # IB23001) by iBlot® 2 Dry Blotting System (Product # IB21001). The blot was probed with Goat anti-Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor 660 (Product # A-21054, 1:2000 dilution) and detected using the iBright FL1500 (Product # A44115). Silver staining was performed to establish equivalent loading of purified proteins using the Pierce™ Silver Stain Kit (Product # 24612) (Fig. b). This antibody also showed reactivity towards Rat IgG light chain.



mTert induction in p21-positive cells counteracts capillary rarefaction and pulmonary emphysema. EMBO Rep (2024)

In vivo identification of Drosophila rhodopsin interaction partners by biotin proximity labeling. Sci Rep (2024)

Tissue-like environments shape functional interactions of HIV-1 with immature dendritic cells. EMBO Rep (2023)

The bacterial effector GarD shields Chlamydia trachomatis inclusions from RNF213-mediated ubiquitylation and destruction. Cell Host Microbe (2022)

Suppressor of Fused Regulation of Hedgehog Signaling is Required for Proper Astrocyte Differentiation. Stem Cells Dev (2022)

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization. Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any Product will conform to such model or sample. NO OTHER WARRANTIES, EXPRESS OR IMPLIED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCT(S) AT SELLER'S SOLE OPTION. THERE IS NO OBLIGATION TO REPAIR, REPLACE OR REFUND FOR PRODUCTS AS THE RESULT OF (i) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (ii) MISUSE, FAULT OR NEGLIGENCE OF OR BY BUYER, (iii) USE OF THE PRODUCTS IN A MANNER FOR WHICH THEY WERE NOT DESIGNED, OR (iv) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressly stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses, or any type of consumption by or application to human or animals.